



Commutative bidifferential algebra

DOI:

[10.1016/j.jalgebra.2022.07.012](https://doi.org/10.1016/j.jalgebra.2022.07.012)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Leon Sanchez, O., & Moosa, R. (2022). Commutative bidifferential algebra. *Journal of Algebra*.
<https://doi.org/10.1016/j.jalgebra.2022.07.012>

Published in:

Journal of Algebra

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



COMMUTATIVE BIDIFFERENTIAL ALGEBRA

OMAR LEÓN SÁNCHEZ AND RAHIM MOOSA

ABSTRACT. Motivated by the Poisson Dixmier-Moeglin equivalence problem, a systematic study of commutative unitary rings equipped with a *biderivation*, namely a binary operation that is a derivation in each argument, is here begun, with an eye toward the geometry of the corresponding *B-varieties*. Foundational results about extending biderivations to localisations, algebraic extensions and transcendental extensions are established. Resolving a deficiency in Poisson algebraic geometry, a theory of base extension is achieved, and it is shown that dominant *B*-morphisms admit generic *B*-fibres. A bidifferential version of the Dixmier-Moeglin equivalence problem is articulated.

CONTENTS

1. Introduction	1
2. Bidifferential algebra	2
3. Bidifferential-algebraic geometry	13
4. The bidifferential Dixmier-Moeglin equivalence problem	17
Appendix A. Extending derivations	21
References	22

1. INTRODUCTION

In this paper we begin a systematic study of commutative *bidifferential* rings; that is, commutative rings equipped with a binary function $\{\cdot, \cdot\} : R \times R \rightarrow R$ which is a derivation in each argument. We are primarily interested in the case when R is a finitely generated integral algebra over a field k – so that we are really talking about affine algebraic varieties equipped with some additional structure, what we call here *B-varieties*¹. In fact, the motivating examples are *Poisson varieties*, namely when $\{\cdot, \cdot\}$ is in addition k -bilinear, skew symmetric, and satisfies the Jacobi identity. But from our point of view, it is more natural to drop these additional assumptions on $\{\cdot, \cdot\}$, at least for now.

In order to illustrate our approach, and the benefit of working beyond the Poisson setting, consider the following question: Given a dominant morphism of affine Poisson varieties $\phi : X \rightarrow Y$ over a field k (of characteristic zero) can we make sense of the generic fibre of ϕ as a Poisson variety? The generic fibre is obtained by taking

Date: July 6, 2022.

2020 Mathematics Subject Classification. 12H05, 13N15, 16W25, 17B63.

Key words and phrases. biderivation, Poisson bracket, D-variety, Dixmier-Moeglin equivalence.

Acknowledgements: R. Moosa was partially supported by an NSERC Discovery Grant.

¹In this terminology, and indeed in our general approach, we are informed by the theory of *D-varieties* as introduced by Buium [6].

a base extension to $L := k(Y)$, namely $\phi_L : X_L \rightarrow Y_L$, and then considering the fibre $\phi_L^{-1}(\alpha)$ where $\alpha \in Y_L(L)$ is a generic point over k . Naively, one might expect that there be canonical Poisson structures on X_L and Y_L , that ϕ_L will be a Poisson morphism, and that $\phi_L^{-1}(\alpha)$ will be a Poisson subvariety of X_L . In fact, all three of these expectations fail separately. One of the main technical accomplishments of this paper (Theorem 3.9 below) is to find a particular bidifferential structure on X_L and Y_L , compatible with the original Poisson structures, such that ϕ_L is a morphism of B -varieties and $\phi_L^{-1}(\alpha)$ is a B -subvariety of X_L . The bidifferential structure on X_L is not canonical in the sense that it depends on ϕ , and not just on the Poisson structure on X and L . Nor is the biderivation we construct necessarily a Poisson bracket. Nevertheless, by working in the more flexible category of B -algebraic geometry we can make good sense of generic fibres of dominant maps.

Our original motivation is the Poisson Dixmier-Moeglin equivalence (PDME) problem, some aspects of which remain open. See, for example, [5, 1, 10]. In our efforts to bring to bear more ideas from differential-algebraic geometry and model theory to this problem, it became clear that fundamental constructions, like base extension and generic fibres, were necessary, and that these in turn required us to move out of the strictly Poisson setting.

From the model theoretic point of view, a central difficulty is that the class of existentially closed bidifferential fields in characteristic zero is not elementary. This is pointed out in §2.3 below. So while our intuition from the model theory of differentially closed fields can still serve as a general guide, all the work has to be done algebraically without recourse to the notions and techniques of geometric stability theory.

Bidifferential algebra has certainly been considered before, but rarely for commutative rings. The preoccupations of the subject as it exists in the literature [3, 4, 8] are rather different than our own, focusing more on classifying and describing biderivations on noncommutative (semi)prime rings, and the results there tend to trivialise in the commutative case.

Here is a plan of the paper. In §2 we develop the foundations of the subject. This includes establishing the basic extension results (to localisations in Theorem 2.8, to algebraic extensions in Theorem 2.10, and to transcendental extension in Theorem 2.13). It also includes the construction of a biderivation on certain tensor products motivated by the generic fibre problem discussed above – this is Theorem 2.19 below. Then in §3 we consider the geometric counterpart of the algebra developed earlier. In particular we introduce B -varieties and show that dominant B -morphisms admit generic B -fibres. Finally, in §4 we articulate the Dixmier-Moeglin Equivalence problem in this setting, as a generalisation of the PDME, and establish some preliminary results (Propositions 4.3, 4.8, 4.9) whose proofs illustrate how commutative bidifferential algebra and the constructions appearing in earlier sections can contribute.

Rings throughout are unital and commutative.

2. BIDIFFERENTIAL ALGEBRA

A *bidifferential ring* is a pair $(R, \{\cdot, \cdot\})$ where R is a (commutative unitary) ring and $\{\cdot, \cdot\} : R \times R \rightarrow R$ is a function such that for each $r \in R$ both $\{r, \cdot\}$ and $\{\cdot, r\}$, called the associated *hamiltonians*, are derivations on R . We call $\{\cdot, \cdot\}$ a *biderivation*.

Examples abound. Every Poisson bracket is a biderivation, and these are the main motivating examples. Another source of examples comes from taking linear combinations of products of derivations: If $\{\delta_1, \dots, \delta_m\}$ and $\{\partial_1, \dots, \partial_m\}$ are derivations on a ring R , and $\alpha_1, \dots, \alpha_m \in R$, then

$$\{r, s\} := \alpha_1 \delta_1(r) \partial_1(s) + \dots + \alpha_m \delta_m(r) \partial_m(s)$$

defines a biderivation on R .

Suppose $(R, \{\cdot, \cdot\})$ is a bidifferential ring. A *bidifferential ideal* of $(R, \{\cdot, \cdot\})$ is an ideal $I \subseteq R$ satisfying $\{R, I\} \subseteq I$ and $\{I, R\} \subseteq I$ – that is, an ideal that is preserved by all the associated hamiltonians. A *morphism* of bidifferential rings, $\phi : (R, \{\cdot, \cdot\}) \rightarrow (S, \{\cdot, \cdot\})$, is a ring homomorphism such that $\phi\{a, b\} = \{\phi a, \phi b\}$ for all $a, b \in R$. The reader can easily verify for themselves the first properties of this category; for example, that the kernel of a morphism of bidifferential rings is a bidifferential ideal, and that the quotient of a bidifferential ring by a bidifferential ideal admits a unique biderivation making the quotient map a morphism of bidifferential rings. But in verifying further properties, especially having to do with extensions of bidifferential rings, it becomes convenient to work in the following more general (if technical) setting.

2.1. Multisorted biderivations. We extend the notion of biderivation so that the arguments accept values from different rings and output values in a module. This will allow us to treat all three sorts individually.

Let \mathcal{K} denote the collection of triples (R, S, M) where R and S are rings and M is an additive group equipped with both an R -module and S -module structure. A *biderivation on (R, S, M)* is then a function $\{\cdot, \cdot\} : R \times S \rightarrow M$ such that

- $\{r, \cdot\} : S \rightarrow M$ is an M -valued derivation on S , for each $r \in R$, and
- $\{\cdot, s\} : R \rightarrow M$ is an M -valued derivation on R , for each $s \in S$.

We call $\{r, \cdot\}$ and $\{\cdot, s\}$ the associated *hamiltonians*. We often say $(R, S, M, \{\cdot, \cdot\})$ is a biderivation to mean that $(R, S, M) \in \mathcal{K}$ and $\{\cdot, \cdot\}$ is a biderivation on (R, S, M) .

We recover (one-sorted) bidifferential rings as biderivations on a triple of the form (R, R, R) , where both module structures (namely of the third sort over the first and over the second) are obtained by viewing R as a module over itself in the natural way. In this case we write $(R, \{\cdot, \cdot\})$ instead of $(R, R, R, \{\cdot, \cdot\})$.

An intermediate level of generalisation that arises frequently is when $R = S$ and the two R -module structures on M agree. In that case we say that $\{\cdot, \cdot\}$ is an *M -valued biderivation on R* . We may write this as $(R, \{\cdot, \cdot\})$ as well, but only when it is clear from context in which module the biderivation takes its values.

By a *morphism in \mathcal{K}* , denoted $\phi : (R, S, M) \rightarrow (R', S', M')$, we mean a triple (ϕ_1, ϕ_2, ϕ_3) such that

- $\phi_1 : R \rightarrow R'$ and $\phi_2 : S \rightarrow S'$ are ring homomorphisms, and
- $\phi_3 : M \rightarrow M'$ is a group homomorphism that is both R -linear and S -linear, in the sense that, for all $x \in M$,
 - $\phi_3(rx) = \phi_1(r)\phi_3(x)$ for all $r \in R$ and
 - $\phi_3(sx) = \phi_2(s)\phi_3(x)$ for all $s \in S$.

Given biderivations on (R, S, M) and (R', S', M') in \mathcal{K} , we say that a morphism $\phi : (R, S, M) \rightarrow (R', S', M')$ is *bidifferential* if

$$\phi_3(\{r, s\}) = \{\phi_1 r, \phi_2 s\}$$

for all $r \in R$ and $s \in S$. We denote this by $\phi : (R, S, M, \{\cdot, \cdot\}) \rightarrow (R', S', M', \{\cdot, \cdot\})$, and say that the biderivation on (R', S', M') *extends* that on (R, S, M) . Note that, whenever we can get away with it, we will use the same symbol to denote the base biderivation and its extension.

The following is the main extension lemma for biderivations.

Proposition 2.1. *Suppose*

- $\phi : (R, S, M) \rightarrow (R', S', M')$ is a morphism in \mathcal{K} with $\phi_1 = \text{id}_R$,
- $\{\cdot, \cdot\}$ is a biderivation on (R, S, M) ,
- $a \in S'$, and
- $\delta : R \rightarrow M'$ is an M' -valued derivation on R .

Suppose each hamiltonian $\{r, \cdot\} : S \rightarrow M$ extends to a unique M' -valued derivation on S' taking a to $\delta(r)$. Then $\{\cdot, \cdot\}$ extends to a unique biderivation on (R, S', M') satisfying $\{a, \cdot\} = \delta$.

Proof. For each $r \in R$, let $\delta_r : S' \rightarrow M'$ denote the (unique) derivation extending $\{r, \cdot\} : S \rightarrow M$ and taking a to $\delta(r)$. Then we have no choice but to define $\{\cdot, \cdot\}$ on (R, S', M') by $\{r, s'\} := \delta_r(s')$. In particular, uniqueness will follow from existence. That this does define an extension of $(R, S, M, \{\cdot, \cdot\})$ follows immediately from the fact that each δ_r extends $\{r, \cdot\} : S \rightarrow M$. So it remains only to show that $(R, S', M', \{\cdot, \cdot\})$ is a biderivation.

Given $r \in R$, we defined $\{r, \cdot\} : S' \rightarrow M'$ to agree with δ_r , and hence these are M' -valued derivations on S' by assumption.

Given $s' \in S'$, we need to show that $\{\cdot, s'\} : R \rightarrow M'$ is a derivation. That is, for all $r_1, r_2 \in R$ we need to show:

$$\begin{aligned} \{r_1 + r_2, s'\} &= \{r_1, s'\} + \{r_2, s'\}, \quad \text{and} \\ \{r_1 r_2, s'\} &= r_2 \{r_1, s'\} + r_1 \{r_2, s'\}. \end{aligned}$$

Fixing $r_1, r_2 \in R$ and letting $s' \in S$ vary, this is equivalent to showing that:

$$\begin{aligned} (1) \quad \delta_{r_1+r_2} &= \delta_{r_1} + \delta_{r_2}, \quad \text{and} \\ (2) \quad \delta_{r_1 r_2} &= r_2 \delta_{r_1} + r_1 \delta_{r_2}, \end{aligned}$$

which is what we now do.

For (1), note that both $\delta_{r_1+r_2}$ and $\delta_{r_1} + \delta_{r_2}$ are M' -valued derivations on S' that extends $\{r_1 + r_2, \cdot\} : S \rightarrow M$. Moreover,

$$\delta_{r_1+r_2}(a) = \delta(r_1 + r_2) = \delta(r_1) + \delta(r_2) = \delta_{r_1}(a) + \delta_{r_2}(a) = (\delta_{r_1} + \delta_{r_2})(a).$$

So, by the uniqueness assumption, we must have $\delta_{r_1+r_2} = \delta_{r_1} + \delta_{r_2}$ on S' .

For (2), note that both $\delta_{r_1 r_2}$ and $r_1 \delta_{r_2} + r_2 \delta_{r_1}$ are M' -valued derivations on S' that extend $\{r_1 r_2, \cdot\} : S \rightarrow M$. Moreover,

$$\delta_{r_1 r_2}(a) = \delta(r_1 r_2) = r_1 \delta(r_2) + r_2 \delta(r_1) = r_1 \delta_{r_2}(a) + r_2 \delta_{r_1}(a) = (r_1 \delta_{r_2} + r_2 \delta_{r_1})(a).$$

Hence $\delta_{r_1 r_2} = r_1 \delta_{r_2} + r_2 \delta_{r_1}$ follows by uniqueness. \square

Remark 2.2. The following observations are worth keeping in mind:

- (a) Proposition 2.1 is often applied with $a = 0$ and $\delta = 0$, in which case it simply says that if all the hamiltonians extend uniquely from $S \rightarrow M$ to $S' \rightarrow M'$ then $\{\cdot, \cdot\}$ extends uniquely from (R, S, M) to (R, S', M') .

- (b) Proposition 2.1 is often applied when S' does not admit any nontrivial derivations that are S -linear – the so called *S -differentially trivial* case. In that case, one has automatically that each hamiltonian has at most one extension to S' (as the difference of two would be S -linear), and the only thing to verify before application is existence.
- (c) Of course, the proposition also works with the roles of R and S reversed – that is, we can keep S fixed and extend R to some R' .

The multi-sorted generalisation of a bidifferential ideal is the following:

Definition 2.3. A *bidifferential ideal pair* of a biderivation $(R, S, M, \{\cdot, \cdot\})$ is a pair of ideals $I \subseteq R$ and $J \subseteq S$ such that $\{I, S\} \subseteq IM$ and $\{R, J\} \subseteq JM$.

One of the complications in bidifferential algebra is that the extension of a bidifferential ideal need not be bidifferential. For example:

Example 2.4. Fix a ring R . It is not hard to construct a biderivation on the polynomial ring in two variables, $R[x, y]$, with the property that $\{\cdot, \cdot\}$ is trivial on $R[x]$ but such that $\{x, y\} = x$. For example, one can use Corollary 2.14 below to verify this. In fact, when $R = k$ is a field, one can do it so that $\{\cdot, \cdot\}$ is a Poisson bracket over k . In any case, we thus have an extension of bidifferential rings; $(R[x], 0) \rightarrow (R[x, y], \{\cdot, \cdot\})$. The ideal $(x-1)R[x]$ is bidifferential in $(R[x], 0)$ as every ideal is, but, since $\{x-1, y\} = \{x, y\} - \{1, y\} = x$, we see that $(x-1)R[x, y]$ is not a bidifferential ideal in the extension $(R[x, y], \{\cdot, \cdot\})$.

We will thus have to restrict our attention to certain extensions of bidifferential rings. Here is the definition in the general multi-sorted setting:

Definition 2.5 (Lifting extensions). An extension of biderivations,

$$\phi : (R, S, M, \{\cdot, \cdot\}) \rightarrow (R', S', M', \{\cdot, \cdot\}),$$

is said to be *lifting* if whenever (I, J) is a bidifferential ideal pair in (R, S, M) then (IR', JS') is a bidifferential ideal pair in (R', S', M') .

We are using the usual commutative algebra conventions here, namely IR' denotes the ideal $\phi_1(I)R'$ of R' generated by the image of I in R' , and similarly for $JS' := \phi_2(J)S'$.

The following is a useful criterion for extensions to be lifting.

Proposition 2.6. *Suppose $\phi : (R, S, M, \{\cdot, \cdot\}) \rightarrow (R', S', M', \{\cdot, \cdot\})$ is a differential morphism with $\phi_1 = \text{id}_R$. If S' differentially trivial² over S then ϕ is lifting.*

Proof. Fix a bidifferential ideal pair (I, J) in (R, S, M) . We need to show that (I, JS') is bidifferential in (R, S', M') .

Let $JM' := (\phi_2(J)S') \cdot M'$, for short. We first show that $\{R, JS'\} \subseteq JM'$. By additivity, it suffices to verify that $\{r, xs'\} \in JM'$ for all $r \in R, x \in J$, and $s' \in S'$. This follows from the following computation:

$$\begin{aligned} \{r, xs'\} &= \{r, \phi_2(x)s'\} \\ &= \phi_2(x)\{r, s'\} + s'\{r, \phi_2(x)\} \\ &= \phi_2(x)\{r, s'\} + s'\phi_3\{r, x\}. \end{aligned}$$

²That is, S' admits no nontrivial S -linear derivations – see Definition A.1

Now, $\phi_2(x)\{r, s'\}$ is visibly in JM' . For the second term, note that, as (I, J) is a bidifferential pair, $\{r, x\} \in JM$, and hence $\phi_3\{r, x\} \in JM'$. So $\{r, xs'\} \in JM'$.

It remains to prove that $\{I, S'\} \subseteq IM'$. This is where differential triviality is used. Fix $x \in I$ and consider the derivation $D := \pi \circ \{x, \cdot\} : S' \rightarrow M'/IM'$. Here $\pi : M' \rightarrow M'/IM'$ is the quotient map. We check that D is S -linear. Indeed, for any $s \in S$, we have

$$\begin{aligned} D(\phi_2(s)) &= \pi\{x, \phi_2(s)\} \\ &= \pi\phi_3\{x, s\} \\ &= 0 \quad \text{as } \{x, s\} \in \{I, S\} \subseteq IM. \end{aligned}$$

So by differential triviality, $D = 0$. But this says precisely that $\{x, S'\} \subseteq IM'$. As $x \in I$ was arbitrary, we have $\{I, S'\} \subseteq IM'$. \square

2.2. Extending bidifferential rings. We can now prove the basic results about extensions of bidifferential rings. We start with observing that differential triviality ensures uniqueness of extensions.

Lemma 2.7. *Suppose R is an A -algebra and M is an R -module. If R admits no nontrivial M -valued A -linear derivations then every M -valued biderivation on A has at most one extension to R .*

Proof. Suppose, toward a contradiction, that $(R, R, M, \{\cdot, \cdot\}_1)$ and $(R, R, M, \{\cdot, \cdot\}_2)$ are distinct extensions of $(A, A, M, \{\cdot, \cdot\})$. Then there exists $r \in R$ such that the hamiltonians $\delta_1 := \{r, \cdot\}_1$ and $\delta_2 := \{r, \cdot\}_2$ are distinct M -valued derivations on R . If δ_1 and δ_2 agreed on A then $\delta_1 - \delta_2$ would be an A -linear M -valued derivation on R , and differential triviality would force $\delta_1 = \delta_2$. Hence, there is $a \in A$ such that $\delta_1(a) \neq \delta_2(a)$. But this means that $d_1 := \{\cdot, a\}_1$ and $d_2 := \{\cdot, a\}_2$ are distinct M -valued derivations on R . But d_1 and d_2 do agree on A because $\{\cdot, \cdot\}_1$ and $\{\cdot, \cdot\}_2$ both extend $\{\cdot, \cdot\}$ and $a \in A$. So $d_1 - d_2$ is a nontrivial A -linear M -valued derivation on R , contradicting our assumption. \square

Theorem 2.8 (Localisation). *Suppose $\{\cdot, \cdot\}$ is an M -valued biderivation on R , and $\Sigma \subseteq R \setminus \{0\}$ is a multiplicatively closed set. Then there is a unique extension of $\{\cdot, \cdot\}$ to an $\Sigma^{-1}M$ -valued biderivation on $\Sigma^{-1}R$. Moreover, this is a lifting extension.*

Proof. By the quotient rule, $\Sigma^{-1}R$ is differentially trivial over R , and hence uniqueness will follow automatically by Lemma 2.7. On the other hand, every M -valued derivation on R does extend to a $\Sigma^{-1}M$ -valued derivation on $\Sigma^{-1}R$. So we can apply Proposition 2.1 with $(a = 0, \delta = 0)$ twice, once to extend from $(R, R, M, \{\cdot, \cdot\})$ to $(R, \Sigma^{-1}R, \Sigma^{-1}M)$, and then to extend to $(\Sigma^{-1}R, \Sigma^{-1}R, \Sigma^{-1}M)$. This is lifting because, by differential triviality, Proposition 2.6 applies to both. \square

Corollary 2.9. *A biderivation on an integral domain extends uniquely to the fraction field.*

Proof. Apply Theorem 2.8 to $M = R$ and $\Sigma = R \setminus \{0\}$. \square

Next we wish to show that we can lift biderivations uniquely to (separably) algebraic extensions. We articulate a rather more general, though somewhat technical, version that allows the base of an algebra to be extended in this way. The greater level of generality is in fact necessary for what we do in the next section.

Theorem 2.10 (Algebraic base extensions). *Suppose $A \subseteq B$ is an extension of integral domains, $b \in B$ is separably algebraic over $\text{Frac}(A)$, and $\{\cdot, \cdot\}$ is a biderivation on an A -algebra R with values in $R \otimes_A B$. Then there is a nonzero $f \in A[b]$ such that $\{\cdot, \cdot\}$ extends uniquely to a biderivation on $R \otimes_A A[b]_f$ with values in $R \otimes_A B_f$. Moreover, this is a lifting extension.*

Proof. Note that in the statement the R -module structure on $R \otimes_A B$, and the $(R \otimes_A A[b])$ -module structure on $R \otimes_A B_f$, are the natural ones.

The first thing we do is reduce to the case when the minimal polynomial of b over $\text{Frac}(A)$ has coefficients in A . Indeed, let $P \in A[t]$ be a nonzero polynomial of minimal degree such that $P(b) = 0$, and let a be the leading coefficient of P . By Theorem 2.8 we have a unique extension of $\{\cdot, \cdot\}$ to the localisation $R_a = R \otimes_A A_a$ with values in $(R \otimes_A B)_a = R_a \otimes_{A_a} B_a$. Now, the minimal polynomial of b over $\text{Frac}(A_a) = \text{Frac}(A)$, namely $\frac{1}{a}P$, does have coefficients in A_a . So, assuming we have proved the proposition in this case, we would get a nonzero $g \in A_a[b]$ and a unique extension of $\{\cdot, \cdot\}$ to $R_a \otimes_{A_a} A_a[b]_g = R \otimes_A A[b]_{ag}$ with values in $R_a \otimes_{A_a} B_{ag} = R \otimes_A B_{ag}$. Moreover this extension would be a lifting extension of the original $\{\cdot, \cdot\}$. Hence $f := ag$ would witness the truth of the proposition.

So we may assume that the minimal polynomial of b over $\text{Frac}(A)$ has coefficients in A . We denote this minimal polynomial by P and set $f := \frac{dP}{dt}(b)$. By separability, f is a nonzero element of $A[b]$. It is pointed out in the appendix, as Fact A.2, that under these conditions every derivation $d : A \rightarrow S$ (where S is any B -algebra) extends uniquely to $d : A[b] \rightarrow S_f$. Combining this with the fact that one can naturally take tensor products of derivations (see Fact A.4), we get that every derivation $d : R \rightarrow R \otimes_A B$ extends uniquely to a derivation $d : R \otimes_A A[b] \rightarrow R \otimes_A B_f$. Now, applying this to each each hamiltonian $\{r, \cdot\}$, viewed as $R \otimes_A B$ -valued, we see that Proposition 2.1 applies (with $a = 0$ and $\delta = 0$). We obtain an extension of $\{\cdot, \cdot\}$ to $(R, R \otimes_A A[b], R \otimes_A B_f)$. Next, applying the same fact to the hamiltonians $\{\cdot, s\}$ for each $s \in R \otimes_A A[b]$, we can invoke Proposition 2.1 again, this time to the first argument, so as to obtain a further extension to $(R \otimes_A A[b], R \otimes_A A[b], R \otimes_A B_f)$. It then extends to $(R \otimes_A A[b]_f, R \otimes_A A[b]_f, R \otimes_A B_f)$ by Theorem 2.8.

Note that if d is an A -linear derivation on $A[b]_f$, with values in any $A[b]_f$ -module M , then $P(b) = 0$ forces $0 = \frac{dP}{dt}(b)db = fdb$, and hence $db = 0$ by multiplying through by $\frac{1}{f}$. So $A[b]_f$ is differentially trivial over A , and it follows that $R \otimes_A A[b]_f$ is differentially trivial over R . By Lemma 2.7, the extension we have built is therefore unique. Also, applying Proposition 2.6 to

$$(R, R, R \otimes_A B, \{\cdot, \cdot\}) \rightarrow (R, R \otimes_A A[b]_f, R \otimes_A B_f, \{\cdot, \cdot\})$$

and then to

$$(R, R \otimes_A A[b]_f, R \otimes_A B_f, \{\cdot, \cdot\}) \rightarrow (R \otimes_A A[b]_f, R \otimes_A A[b]_f, R \otimes_A B_f, \{\cdot, \cdot\})$$

shows that the extension we have built is lifting. \square

Remark 2.11. The proposition will only be used when $R \otimes_A B$ has no A -torsion. It is not hard to show that in that case, in order to extend the hamiltonions to $R \otimes_A A[b]$, one does not require that the minimal polynomial of b over $\text{Frac}(A)$ has coefficients in A . (See the proof of Fact A.2.) Hence, assuming that $R \otimes_A B$ has no A -torsion, we can, in Theorem 2.10, take f to be $\frac{dP}{dt}(b)$ where $P \in A[t]$ is a polynomial of minimal degree such that $P(b) = 0$.

Corollary 2.12. *A biderivation on a field extends uniquely to the separable closure.*

Proof. This can be deduced from recursive applications of Theorem 2.10, starting with $A = R =: k$ a field, and $B = k^{\text{sep}}$, and extending the biderivation one element at a time. Or, and this may be easier, we can directly apply Proposition 2.1 to first extend $\{\cdot, \cdot\}$ uniquely to $(k, k^{\text{sep}}, k^{\text{sep}})$, and then apply Proposition 2.1 again to extending the first argument to k^{sep} . Both stages use the fact that derivations $d : k \rightarrow k^{\text{sep}}$ extend uniquely to a derivation on k^{sep} . \square

Theorem 2.13 (Transcendental base extensions). *Suppose $A \subseteq B$ is an extension of rings and $b \in B$ is transcendental over A . Suppose R is an A -algebra, $D : R \rightarrow R \otimes_A B$ is a derivation, and $E : R \otimes_A A[b] \rightarrow R \otimes_A B$ is a derivation. Every $(R \otimes_A B)$ -valued biderivation on R extends uniquely to a $(R \otimes_A B)$ -valued biderivation on $R \otimes_A A[b]$ satisfying*

- $\{r \otimes 1, 1 \otimes b\} = D(r)$ for all $r \in R$, and
- $\{1 \otimes b, x\} = E(x)$ for all $x \in R \otimes_A A[b]$.

Proof. First apply Proposition 2.1 with $S = R$, $M = M' = R \otimes_A B$, $S' = R \otimes_A A[b]$, and with $a := 1 \otimes b$ and $\delta := D$. Note that $R \otimes_A A[b] = R[a]$ is isomorphic to the polynomial ring in one variable over R , and so for each $r \in R$ it is true that $\{r, \cdot\} : R \rightarrow R \otimes_A B$ does have a unique extension to a derivation on $R[a]$ taking a to $\delta(r)$. Hence, we get a unique extension $(R, R \otimes_A A[b], R \otimes_A B, \{\cdot, \cdot\})$. Now we apply Proposition 2.1 again, this time to extend the first argument from R to $R \otimes_A A[b]$. For that, we use again $a := 1 \otimes b$ and this time $\delta := E$. For each $x \in R \otimes_A A[b]$, the hamiltonian $\{\cdot, x\} : R \rightarrow R \otimes_A B$ has a unique extension to a derivation on $R[a] = R \otimes_A A[b]$ taking a to $\delta(x)$. The extended $(R \otimes_A B)$ -valued biderivation on $R \otimes_A A[b]$ that we obtain has by construction the two desired properties, and is by construction unique such. \square

Corollary 2.14. *Every biderivation on R extends to the polynomial ring $R[t]$. Moreover, given a pair of derivation $D : R \rightarrow R[t]$ and $E : R[t] \rightarrow R[t]$, there is a unique extension of $\{\cdot, \cdot\}$ to $R[t]$ satisfying $\{\cdot, t\} \upharpoonright_R = D$ and $\{t, \cdot\} = E$.*

Proof. Apply Theorem 2.13 with $A = R$, $B = R[t]$, $b = t$, and D, E as given. \square

2.3. A model-theoretic aside. We can view bidifferential rings as first-order structures in the language L of rings augmented by a binary function symbol for the biderivation. But there is no good model theory of biderivations in the sense that the class of existentially closed bidifferential fields is not elementary. Let us show how this follows from some of the basic extension results proved above.

Proposition 2.15. *The L -theory of bidifferential fields of characteristic zero does not admit a model companion.*

Proof. First of all, any biderivation extends (uniquely) to the separable closure by Corollary 2.12. It follows that an existentially closed bidifferential field of characteristic zero is algebraically closed.

By the *constants* of a bidifferential ring $(R, \{\cdot, \cdot\})$ we mean the subring of elements $r \in R$ such that the associated hamiltonians $\{r, \cdot\}$ and $\{\cdot, r\}$ are identically zero. We observe that the constant subfield of an existentially closed bidifferential field $(F, \{\cdot, \cdot\})$ is always \mathbb{Q}^{alg} . Indeed, consider the function field $F(t)$ in one variable.

Given any derivation $D : F \rightarrow F$, we can, using Corollaries 2.9 and 2.14, build a biderivation on $F(t)$ extending $\{\cdot, \cdot\}$ on F and such that $\{\cdot, t\} \upharpoonright_F = D$. Suppose, now, that we are given $r \in F \setminus \mathbb{Q}^{\text{alg}}$. Then we can find D such that $D(r) \neq 0$. Hence $\{r, t\} = D(r) \neq 0$. It follows by existential closedness, that there is some $s \in F$ such that $\{r, s\} \neq 0$. So r is not a constant in $(F, \{\cdot, \cdot\})$.

Since the constants, which form a 0-definable infinite set, does not grow within the class of existentially closed bidifferential fields, that class cannot be elementary. \square

2.4. Tensor products. Unlike in the differential case (see Fact A.4), biderivations are not always compatible with tensor products. The following example shows that we cannot always equip the tensor product of bidifferential rings with a biderivation:

Example 2.16. Let k be a field and consider $R := k[x, y] \otimes_{k[x]} k$, where the polynomial ring $k[x, y]$ is viewed as a $k[x]$ -algebra in the natural way, and k is viewed as a $k[x]$ -algebra via the evaluation at zero homomorphism. Now equip k and $k[x]$ with trivial biderivations, while equipping $k[x, y]$ with the biderivation extending the trivial one on $k[x]$ and satisfying $\{\cdot, y\} \upharpoonright_{k[x]} = \frac{d}{dx}$. One can use Corollary 2.14 to see this exists. We **cannot** equip R with a bidifferential structure extending $(k[x, y], \{\cdot, \cdot\})$. Indeed, if we could, then using that $1 \otimes x = 0$ in R , we would have $0 = \{y \otimes 1, 1 \otimes x\} = \{y \otimes 1, x \otimes 1\} = \{x, y\} \otimes 1 = 1 \otimes 1$, which is a contradiction as R is not the trivial ring.

Even if we are interested primarily in tensor products over fields, as we are, problems persist. Let us fix a bidifferential field $(k, \{\cdot, \cdot\})$ and consider the category of *bidifferential k -algebras*, namely k -algebras equipped with a biderivation that extends $\{\cdot, \cdot\}$ on k . We would like to equip the tensor product $R \otimes_k S$ of bidifferential k -algebras with a suitable biderivation. But, again unlike for differential rings, there is no truly canonical choice. At first glance, at least when the biderivation is trivial on k , it does seem that a canonical choice would be to define

$$(3) \quad \{r_1 \otimes s_1, r_2 \otimes s_2\} = \{r_1, r_2\} \otimes s_1 s_2 + r_1 r_2 \otimes \{s_1, s_2\},$$

as is done in Poisson algebra. Indeed, (3) induces the unique biderivation on $R \otimes_k S$ that extends the biderivations on R and S and that sets $\{r \otimes 1, 1 \otimes s\} = 0$. But, as the following example shows, that choice has some pathological implications.

Example 2.17. Suppose $\{\cdot, \cdot\}$ is trivial on k and make $R = k[x, y]$ the bidifferential k -algebra where $\{x, y\} = x$. (So this is the same as Example 2.4.) If we equip $R \otimes_k R = k[x \otimes 1, y \otimes 1, 1 \otimes x, 1 \otimes y]$, with the canonical biderivation given by (3) then the diagonal ideal $I := (x \otimes 1 - 1 \otimes x, y \otimes 1 - 1 \otimes y)$ is not bidifferential. Indeed, $\{x \otimes 1 - 1 \otimes x, y \otimes 1\} = \{x, y\} \otimes 1 = x \otimes 1$, which is not an element of I .

In any case, when $\{\cdot, \cdot\}$ is not trivial on k , (3) is simply not available to us: if $r, s \in k$ then $\{r \otimes 1, 1 \otimes s\} = \{r \otimes 1, s \otimes 1\}$ which should agree with $\{r, s\} \otimes 1$ and hence $\{r \otimes 1, 1 \otimes s\}$ should *not* be zero if $\{r, s\} \neq 0$ in k .

We propose an alternative bidifferential structure on certain tensor products of bidifferential k -algebras. First, we will mostly be interested in the affine case:

Definition 2.18. An *affine bidifferential k -algebra* is a finitely generated k -algebra that is an integral domain and is equipped with a biderivation extending $(k, \{\cdot, \cdot\})$.

The following statement may seem technical, but it has a clear geometric formulation and motivation that will be expressed later by Theorem 3.6. For now, let us only point out that ι corresponds to a dominant morphism of affine algebraic varieties, and the defining ideal of ι in the tensor product corresponds to the graph of that morphism.

Theorem 2.19. *Suppose $(k, \{\cdot, \cdot\})$ is a bidifferential field, $(R, \{\cdot, \cdot\})$ and $(S, \{\cdot, \cdot\})$ are affine bidifferential k -algebras with $\text{Frac}(S)$ separable over k . Suppose $\iota : S \rightarrow R$ is a k -linear bidifferential embedding. Then there exists a nonzero $f \in S$ and a biderivation on $R \otimes_k S_f$ satisfying the following properties:*

- (a) $(R \otimes_k S_f, \{\cdot, \cdot\})$ is a lifting extension of $(R, \{\cdot, \cdot\})$.
- (b) $(R \otimes_k S_f, \{\cdot, \cdot\})$ is an extension of $(S, \{\cdot, \cdot\})$.
- (c) the subring $\iota(S) \otimes_k S_f \subseteq R \otimes_k S_f$ is bidifferential, and it is a lifting extension of $(\iota(S), \{\cdot, \cdot\})$.
- (d) The defining ideal of ι , namely the ideal generated by $\iota s \otimes 1 - 1 \otimes s$ for all $s \in S$, is a bidifferential ideal of $(R \otimes_k S_f, \{\cdot, \cdot\})$.

Remark 2.20. In particular, if we take $S = R$ and $\iota = \text{id}$ then this gives us a biderivation on (a localisation) of $R \otimes_k R$ avoiding the pathologies exhibited by Example 2.17.

Proof of Theorem 2.19. By the separable form of Noether's normalisation lemma (see [7, 16.18]) there is $y := (y_1, \dots, y_n)$ in S algebraically independent over k such that $\text{Frac}(S)$ is separably algebraic over $k(y)$.

Step 1 of the construction. We extend $\{\cdot, \cdot\}$ from R to an $R \otimes_k S$ -valued biderivation on $R \otimes_k k[y]$ using Theorem 2.13 recursively. For this, we view the biderivation on R as being $R \otimes_k S$ -valued, and the biderivation on S restricting to an $R \otimes_k S$ -valued biderivation on $k[y]$. That is, we have

$$\begin{aligned} \{\cdot, \cdot\} & : R \times R \rightarrow R \otimes_k S \\ \{\cdot, \cdot\} & : k[y] \times k[y] \rightarrow R \otimes_k S \end{aligned}$$

Suppose we have extended these to $R \otimes_k k[y_1, \dots, y_{i-1}]$. Define the derivation $D : R \otimes_k k[y_1, \dots, y_{i-1}] \rightarrow R \otimes_k S$ to be the derivation determined by

$$\begin{aligned} D(r \otimes 1) & = \{r, \iota y_i\} \otimes 1 \text{ for all } r \in R, \text{ and} \\ D(1 \otimes y_j) & = 1 \otimes \{y_j, y_i\} \text{ for all } j < i, \end{aligned}$$

and define $E : R \otimes_k k[y_1, \dots, y_i] \rightarrow R \otimes_k S$ to be the derivation determined by:

$$\begin{aligned} E(r \otimes 1) & = \{\iota y_i, r\} \otimes 1 \text{ for all } r \in R, \text{ and} \\ E(1 \otimes y_j) & = 1 \otimes \{y_i, y_j\} \text{ for all } j \leq i. \end{aligned}$$

Now apply Theorem 2.13 with this D and E to obtain $\{\cdot, \cdot\}$ on $R \otimes_k k[y_1, \dots, y_i]$. Recursively, we obtain a unique $R \otimes_k S$ -valued biderivation on $R \otimes_k k[y]$ that extends the given biderivations on R and $k[y]$ and satisfies:

$$\begin{aligned} (4) \quad \{r \otimes 1, 1 \otimes y_i\} & = \{r, \iota y_i\} \otimes 1 \\ (5) \quad \{1 \otimes y_i, r \otimes 1\} & = \{\iota y_i, r\} \otimes 1 \end{aligned}$$

for all $r \in R$ and $i = 1, \dots, n$.

Step 2 of the construction. We now extend $\{\cdot, \cdot\}$ from $R \otimes_k k[y]$ to $R \otimes_k S$, at the expense of localising S . As S is a finitely generated and $\text{Frac}(S)$ is separably algebraic over $k(y)$, we can write $S = k[y, b]$ where $b = (b_1, \dots, b_m)$ are separably

algebraic over $k(y)$. For each i let $P_i \in k[y, b_1, \dots, b_{i-1}][t]$ be of minimal degree such that $P_i(b_i) = 0$. Let $f := \prod_{i=1}^m \frac{dP_i}{dt}(b_i)$. Applying Theorem 2.10 repeatedly we obtain an extension of $(R \otimes_k k[y], \{\cdot, \cdot\})$ to $(R \otimes_k S_f, \{\cdot, \cdot\})$. Actually, we are using here that $R \otimes_k S$ has no S -torsion to extract from Theorem 2.10 that this particular f works – otherwise we only know there is some f which does. See Remark 2.11.

We now verify (a) through (d).

(a). By construction, $(R, \{\cdot, \cdot\}) \rightarrow (R \otimes_k k[y], \{\cdot, \cdot\}) \rightarrow (R \otimes_k S_f, \{\cdot, \cdot\})$ are extensions. Here we view all the above biderivations as $R \otimes_k S_f$ -valued under the natural identifications. We verify that both steps are lifting extensions. The second is lifting because it is obtained by repeated applications of Theorem 2.10 and then an application of Theorem 2.8, and both of these theorems tell us that the extensions they produce are lifting. To verify that the first extension is lifting, fix a bidifferential ideal pair (I, J) in $(R, \{\cdot, \cdot\})$. We first show that

$$\{I(R \otimes_k k[y]), R \otimes_k k[y]\} \subseteq I(R \otimes_k S).$$

Let $x \in I$. It suffices to show that $\{x \otimes 1, r \otimes 1\}$ and $\{x \otimes 1, 1 \otimes y_i\}$ are in $I(R \otimes_k S)$ for all $r \in R$ and $i = 1, \dots, n$. The former is because $\{x \otimes 1, r \otimes 1\} = \{x, r\} \otimes 1$ and $\{I, R\} \subseteq I$. The second is because $\{x \otimes 1, 1 \otimes y_i\} = \{x, \iota y_i\} \otimes 1$ by (4) and again using that $\{I, R\} \subseteq I$. A similar argument, using (5) and the fact that $\{R, J\} \subseteq J$, shows that $\{R \otimes_k k[y], J(R \otimes_k k[y])\} \subseteq J(R \otimes_k S)$. So $(I(R \otimes_k k[y]), J(R \otimes_k k[y]))$ is a bidifferential ideal pair, as desired.

(b). Let $\{\cdot, \cdot\}'$ be the $(R \otimes_k S_f)$ -valued biderivation on S obtained by restricting $(R \otimes_k S_f, \{\cdot, \cdot\})$ to $S = 1 \otimes S$. We want to show that $\{\cdot, \cdot\}' = \{\cdot, \cdot\}$. Note that $\{\cdot, \cdot\}'$ and $\{\cdot, \cdot\}$ agree on $k[y]$ because by construction,

$$(k[y], \{\cdot, \cdot\}) \rightarrow (R \otimes_k k[y], \{\cdot, \cdot\}) \rightarrow (R \otimes_k S_f, \{\cdot, \cdot\})$$

are extensions. Now, even though $\text{Frac}(S)$ is separably algebraic over $k(y)$, it is not exactly the case that S is differentially trivial over $k[y]$. The problem arises when one considers derivations on S with values in an S -module that has torsion. However, it is true that if M is a torsion-free S -module then S has no nontrivial $k[y]$ -linear M -valued derivations. It follows by Lemma 2.7 that there is at most one M -valued biderivation on S extending any given M -valued biderivation on $k[y]$. Applying this to $M := R \otimes_k S_f$, which has no S -torsion as it is free over S_f , we get that $\{\cdot, \cdot\}' = \{\cdot, \cdot\}$ on S , as desired.

(c). Since $\iota : S \rightarrow R$ is bidifferential and k -linear, $\iota(S)$ is a bidifferential k -subalgebra of R . Inspecting the construction of $\{\cdot, \cdot\}$ on $R \otimes_k S_f$ readily reveals that $\iota(S) \otimes_k S_f$ is a bidifferential subring of $R \otimes_k S_f$. In fact, the restriction of $\{\cdot, \cdot\}$ to $\iota(S) \otimes_k S_f$ is just the biderivation our construction yields if we had viewed ι as $\iota(S)$ -valued rather than as R -valued. So part (a) of this theorem applied to the case when $R = \iota(S)$, gives us that $\iota(S) \otimes_k S_f$ is a lifting extension of $\iota(S)$.

(d). Let $d : S \rightarrow R \otimes_k S_f$ be the function $d(s) := \iota s \otimes 1 - 1 \otimes s$, and let I be the ideal of $R \otimes_k S_f$ generated by $\{d(s) : s \in S\}$. We wish to show that I is a bidifferential ideal. We begin with the following approximation:

Claim 2.21. *Let I_0 be the ideal in $R \otimes_k k[y]$ generated by $\{d(s) : s \in k[y]\}$. Then $I_0(R \otimes_k S_f)$ is a bidifferential ideal.*

Proof of Claim. By construction, and Theorems 2.8 and 2.10, $(R \otimes_k S_f, \{\cdot, \cdot\})$ is a lifting extension of $(R \otimes_k k[y], \{\cdot, \cdot\})$. Hence it suffices to show that I_0 is a

bidifferential ideal. We have to show that $\{x, d(s)\}$ and $\{d(s), x\}$ are in I_0 for all $x \in R \otimes_k k[y]$ and $s \in k[y]$. By symmetry, it will suffice to show the former. It is not hard to see that it suffices to prove this when $x = r \otimes y_i$ and $s = y_j$, for some $r \in R$ and $i, j \leq m$. This is what we do.

We compute

$$\begin{aligned} \{r \otimes y_i, 1 \otimes y_j\} &= \{r \otimes 1, 1 \otimes y_j\}1 \otimes y_i + r \otimes 1\{1 \otimes y_i, 1 \otimes y_j\} \\ &= \{r, \iota y_j\} \otimes y_i + r \otimes \{y_i, y_j\} \quad \text{by (4)} \end{aligned}$$

and

$$\begin{aligned} \{r \otimes y_i, \iota y_j \otimes 1\} &= \{r \otimes 1, \iota y_j \otimes 1\}1 \otimes y_i + r \otimes 1\{1 \otimes y_i, \iota y_j \otimes 1\} \\ &= \{r, \iota y_j\} \otimes y_i + r\{\iota y_i, \iota y_j\} \otimes 1 \quad \text{by (5)} \\ &= \{r, \iota y_j\} \otimes y_i + r\iota(y_i, y_j) \otimes 1 \quad \text{as } \iota \text{ is bidifferential.} \end{aligned}$$

Hence, we get

$$\begin{aligned} \{r \otimes y_i, d(y_j)\} &= \{r \otimes y_i, \iota y_j \otimes 1\} - \{r \otimes y_i, 1 \otimes y_j\} \\ &= r \otimes 1(\iota(\{y_i, y_j\}) \otimes 1 - 1 \otimes \{y_i, y_j\}) \\ &= (r \otimes 1)d(\{y_i, y_j\}) \\ &\in I_0 \end{aligned}$$

as desired. \dashv

To pass from I_0 to I we will use the following derivative-like property of d .

Claim 2.22. *Suppose $P \in S[t]$ and $s \in S$. Then $d(P(s)) = ad(s) + P^d(\iota s \otimes 1)$ where $a \equiv 1 \otimes \frac{dP}{dt}(s) \pmod{I}$. Here P^d is the polynomial over $R \otimes_k S_f$ obtained from P by applying d to the coefficients.*

Proof of Claim. By additivity it suffices to prove this in the case that $P(t) = bt^n$ is a monomial. First, one verifies readily that, for all $s_1, s_2 \in S$,

$$d(s_1 s_2) = d(s_1)(\iota s_2 \otimes 1) + (1 \otimes s_1)d(s_2).$$

From this, by induction on $n > 0$, it follows that

$$d(s^n) = (\iota s^{n-1} \otimes 1 + \iota s^{n-2} \otimes s + \cdots + \iota s \otimes s^{n-2} + 1 \otimes s^{n-1})d(s)$$

and hence

$$\begin{aligned} d(P(s)) &= d(bs^n) \\ &= d(b)(\iota s^n \otimes 1) + (1 \otimes b)(\iota s^{n-1} \otimes 1 + \iota s^{n-2} \otimes s + \cdots + 1 \otimes s^{n-1})d(s) \\ &= P^d(\iota s \otimes 1) + ad(s) \end{aligned}$$

where $a := (1 \otimes b)(\iota s^{n-1} \otimes 1 + \iota s^{n-2} \otimes s + \cdots + 1 \otimes s^{n-1})$. Now, note that

$$\iota s^{n-i} \otimes s^{i-1} \equiv 1 \otimes s^{n-1} \pmod{I}$$

for each $i = 1, \dots, n$. And so,

$$\begin{aligned} a &= (1 \otimes b)(\iota s^{n-1} \otimes 1 + \iota s^{n-2} \otimes s + \cdots + 1 \otimes s^{n-1}) \\ &\equiv (1 \otimes b)(1 \otimes ns^{n-1}) \pmod{I} \\ &= 1 \otimes \frac{dP}{dt}(s) \pmod{I} \end{aligned}$$

as desired. \dashv

Now, to prove that I is bidifferential it suffices to check that $\{x, d(s)\}$ and $\{d(s), x\}$ are in I for all $s \in S$ and $x \in R \otimes_k S_f$. We will prove the former, the latter follows by symmetry.

Recall that S is separably algebraic over $k(y)$. Let $P \in k[y][t]$ be of minimal degree such that $P(s) = 0$. Then

$$\begin{aligned} 0 &= \{x, d(P(s))\} \\ &= \{x, ad(s) + P^d(\iota s \otimes 1)\} \quad \text{by Claim 2.22} \\ &= a\{x, d(s)\} + d(s)\{x, a\} + \{x, P^d(\iota s \otimes 1)\} \\ &\equiv (1 \otimes \frac{dP}{dt}(s))\{x, d(s)\} + \{x, P^d(\iota s \otimes 1)\} \quad \text{mod } I. \end{aligned}$$

Now, as the coefficients of P are in $k[y]$, we have that $P^d(\iota s \otimes 1) \in I_0$. Hence, by Claim 2.21, $\{x, P^d(\iota s \otimes 1)\} \in I_0(R \otimes_k S_f) \subseteq I$. It follows that

$$(1 \otimes \frac{dP}{dt}(s))\{x, d(s)\} \in I.$$

This is a good time to observe that I is (the localisation of) the kernel of the k -linear homomorphism $R \otimes_k S \rightarrow R$ given by $r \otimes s \mapsto r\iota(s)$. In particular, as $\frac{dP}{dt}(s) \neq 0$ by minimal choice of degree, and ι is injective, we have that $(1 \otimes \frac{dP}{dt}(s)) \notin I$. But, as R is an integral domain, I is prime. Hence $\{x, d(s)\} \in I$, as desired. \square

3. BIDIFFERENTIAL-ALGEBRAIC GEOMETRY

In this section, we fix a bidifferential field $(k, \{\cdot, \cdot\})$ of characteristic zero. The following is the bidifferential analogue of the D -varieties studied in differential-algebraic geometry.

Definition 3.1. A B -variety over k is an affine algebraic variety X over k together with a biderivation on $k[X]$ extending $(k, \{\cdot, \cdot\})$.

A B -subvariety of X is a subvariety whose vanishing ideal is a bidifferential ideal.

A B -point of X over k is a k -point which as a singleton is a B -subvariety. The set of all B -points is denoted by $X^\sharp(k)$.

A *morphism* of B -varieties (or a B -morphism) is a regular map $\phi : X \rightarrow Y$ such that the induced k -algebra homomorphism $\phi^* : k[Y] \rightarrow k[X]$ is bidifferential. We say ϕ is a *lifting* morphism if ϕ^* is a lifting extension.

Note that if $\phi : X \rightarrow Y$ is a lifting morphism then $\phi^{-1}(Z)$ is a B -subvariety of X , for every B -subvariety $Z \subseteq Y$. This is not necessarily true if ϕ is not lifting.

The irreducible components of a B -subvariety are themselves B -subvarieties. Indeed, one applies to each of the hamiltonians of $(k[X], \{\cdot, \cdot\})$ the fact that the minimal prime ideals containing a radical differential ideal in a noetherian differential ring are all differential. See [9, §1.2], for instance.

Remark 3.2. It is tempting to associate to a B -variety X the various D -variety structures arising from the hamiltonians. The problem is that there is no reason why, for a given $a \in k[X]$, the derivation $\{a, \cdot\} : k[X] \rightarrow k[X]$ should extend a (k -valued) derivation on the field k , in general it is only $k[X]$ -valued. However, if it happens that $\{a, \cdot\}$ restricts to a derivation $\delta_a : k \rightarrow k$ then one can consider the D -variety (X, s_a) over (k, δ_a) given by the section $s_a : X \rightarrow \tau X$ induced by $\{a, \cdot\}$. (Here τX denotes the prolongation of X with respect to δ_a .) In particular, if

$\{a, \cdot\}$ is k -linear then $s_a : X \rightarrow TX$ puts on X the structure of a D -variety over the constants. For example, if X is a Poisson variety over k , and so all the hamiltonians are k -linear, we can associate to X the collection $\{s_a : X \rightarrow TX : a \in k[X]\}$ of D -variety structures on X induced by the hamiltonians, and these D -variety structures do to some extent capture the Poisson structure. This connection between Poisson and D -variety structures has been exploited in [1, 2, 10] to answer questions around the Poisson Dixmier-Moeglin equivalence.

Given a B -variety X over k and a field extension $k \subseteq L$, it will of course be necessary to sometimes view X as “being over L ”, namely to take the base extension $X_L := X \times_k L$. This suggests questions about the existence and uniqueness of B -variety structures on X_L extending the given one on X . Actually we will only be interested in “compatible” B -variety structures:

Definition 3.3 (Compatible base extension). Suppose $k \subseteq L$ is an extension of bidifferential fields and X is a B -variety over k . By a *compatible* B -variety structure on X_L we mean a biderivation on $L[X_L] = k[X] \otimes_k L$ that extends the given biderivations on $k[X]$ and L , and such that $k[X] \subseteq L[X_L]$ is a lifting extension.

The point is that if we equip X_L with a compatible B -variety structure then whenever Z is a B -subvariety of X , Z_L is a B -subvariety of X_L . As one might by now expect, X_L may not admit a compatible B -variety structure, and even if it did there may be no canonical choice. The following is the geometric formulation of Example 2.4:

Example 3.4. Let X be the affine line \mathbb{A}^1 over k , and let L be the rational function field $k(y)$. So X_L is the “generic fibre” of the co-ordinate projection $\mathbb{A}^2 \rightarrow \mathbb{A}^1$. We put on k and L the trivial biderivation, and on X the trivial B -variety structure. What about $L[X_L] = k(y)[x]$? We can of course equip it too with the trivial biderivation, making X_L a compatible base extension of X . But we could also consider the biderivation on $L[X_L]$ that is trivial on $k(y)$ and $k[x]$ but where $\{x, y\} = x$. Then the corresponding B -variety structure on X_L is not a compatible base extension of X . Indeed, $Z = \{1\}$ is a B -subvariety of X but Z_L is not a B -subvariety of X_L , as $(x-1)k(y)[x]$ is not a bidifferential ideal in $k(y)[x]$ since $\{x-1, y\} = x$.

Fortunately these issues do not arise when taking algebraic base extensions.

Proposition 3.5. *Suppose X is a B -variety over k . Then there is unique B -variety structure on $X_{k^{\text{alg}}}$ extending that on X , and this is a compatible base extension.*

Proof. First, as we are in characteristic zero, there is, by Corollary 2.12, a unique extension of $\{\cdot, \cdot\}$ from k to k^{alg} , which we also denote by $\{\cdot, \cdot\}$. It is with respect to $(k, \{\cdot, \cdot\}) \subseteq (k^{\text{alg}}, \{\cdot, \cdot\})$ that a compatible base extension of X is being claimed.

View the biderivation on $k[X]$ as having values in $k[X] \otimes_k k^{\text{alg}} = k^{\text{alg}}[X_{k^{\text{alg}}}]$. Fix $b \in k^{\text{alg}}$. By Theorem 2.10, applied to $A = k$, $B = k^{\text{alg}}$ and $R = k[X]$, there is an extension of $\{\cdot, \cdot\}$ from $k[X]$ to $k[X] \otimes_k k(b)$, and moreover this is a lifting extension. Now work over $k(b)$ and iterate. By Zorn’s lemma this eventually produces a lifting extension $(k[X], \{\cdot, \cdot\}) \subseteq (k[X] \otimes_k k^{\text{alg}}, \{\cdot, \cdot\}')$. To see that it extends $(k^{\text{alg}}, \{\cdot, \cdot\})$ as well, note that $\{\cdot, \cdot\}'$ restricts to a $(k[X] \otimes_k k^{\text{alg}})$ -valued biderivation on k^{alg} extending $(k, \{\cdot, \cdot\})$. The fact that k^{alg} is differentially trivial over k then implies, by Lemma 2.7, that $\{\cdot, \cdot\}'$ agrees with $\{\cdot, \cdot\}$ on k^{alg} .

So $(k[X] \otimes_k k^{\text{alg}}, \{\cdot, \cdot\}')$ gives $X_{k^{\text{alg}}}$ a B -variety structure that is a compatible base extension of X . Uniqueness follows from Lemma 2.7 using the fact that $k[X] \otimes_k k^{\text{alg}}$ is differentially trivial over $k[X]$. \square

Here is the geometric interpretation of Theorem 2.19.

Theorem 3.6. *Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of irreducible B -varieties over k . Then there exists a nonempty Zariski open subset $Y_0 \subseteq Y$ and a B -variety structure on $X \times Y_0$ such that*

- (a) *the co-ordinate projection $X \times Y_0 \rightarrow X$ is a lifting B -morphism,*
- (b) *the co-ordinate projection $X \times Y_0 \rightarrow Y_0$ is a B -morphism,*
- (c) *$\phi \times \text{id} : X \times Y_0 \rightarrow Y \times Y_0$ induces a B -variety structure on $Y \times Y_0$ such that the co-ordinate projection $Y \times Y_0 \rightarrow Y$ is a lifting B -morphism, and*
- (d) *the graph of $\phi|_{Y_0}$ is a B -subvariety of $X \times Y_0$.*

Proof. Let $R := k[X]$, $S := k[Y]$, and $\iota := \phi^* : k[Y] \rightarrow k[X]$ which is an embedding as ϕ is dominant and is bidifferential as ϕ is a B -morphism. Applying Theorem 2.19 yields a nonzero $f \in k[Y]$ and a biderivation on $k[X] \otimes_k k[Y]_f$, satisfying certain properties. Now, $k[X] \otimes_k k[Y]_f = k[X \times Y_0]$ where $Y_0 := Y \setminus V(f)$. This makes $X \times Y_0$ a B -variety. Now the statements (a) through (d) of the current theorem are just the geometric meanings of the corresponding statements (a) through (d) of Theorem 2.19. \square

3.1. Generic fibres and a universal base extension. With a bit more work we can use Theorem 3.6 to find generic fibres of dominant B -morphisms.

Definition 3.7. Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of irreducible B -varieties over k . By a *generic B -fibre* of ϕ we mean

- a bidifferential field extension $L \supseteq k$,
- compatible base extensions X_L and Y_L such that $\phi_L : X_L \rightarrow Y_L$ is a B -morphism, and
- a B -point $\alpha \in Y_L^\sharp(L)$ that is (Zariski) generic in Y_L over k such that $\phi_L^{-1}(\alpha)$ is a B -subvariety of X_L

We sometimes suppress the data and simply call $X_\alpha := \phi_L^{-1}(\alpha)$ a generic B -fibre over L .

Note, first of all, that fibres over B -points need not be B -subvarieties in general:

Example 3.8. In Example 2.4 (again!) we considered the trivial biderivation on $k[x]$ extended to the biderivation on $k[x, y]$ where $\{x, y\} = x$. So we have the trivial B -variety structure on \mathbb{A}^1 and a B -variety structure on \mathbb{A}^2 such that the first co-ordinate projection $\pi : \mathbb{A}^2 \rightarrow \mathbb{A}^1$ is a B -morphism. Now $1 \in \mathbb{A}^1(k)$ is a B -point as every k -point is. But $\pi^{-1}(1)$ has vanishing ideal $(x-1)k[x, y]$, which we have seen is not a bidifferential ideal.

Theorem 3.9. *Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of irreducible B -varieties over k . Then a generic B -fibre of ϕ exists over $L = k(Y)$.*

Proof. Apply Theorem 3.6 to get a nonempty Zariski open subset $Y_0 \subseteq Y$ and a B -variety structure on $X \times Y_0$ such that the co-ordinate projection to X is a lifting B -morphism. At the same time, we get a B -variety structure on $Y \times Y_0$ such that $\phi \times \text{id} : X \times Y_0 \rightarrow Y \times Y_0$ is a B -morphism and $Y \times Y_0 \rightarrow Y$ is a lifting B -morphism.

Let $L := k(Y) = k(Y_0)$. As localisation is lifting (Theorem 2.8), we have that $k[X] \otimes_k k[Y_0] \subseteq k[X] \otimes_k L = L[X_L]$ and $k[Y] \otimes_k k[Y_0] \subseteq k[Y] \otimes_k L = L[Y_L]$ are lifting extension. In diagrams, we now have:

$$\begin{array}{ccc}
 L[Y_L] & \xrightarrow{\phi_L^*} & L[X_L] \\
 \uparrow & & \uparrow \\
 k[Y] \otimes_k k[Y_0] & \xrightarrow{(\phi \times \text{id})^*} & k[X] \otimes_k k[Y_0] \\
 \uparrow & & \uparrow \\
 k[Y] & \xrightarrow{\phi^*} & k[X]
 \end{array}$$

with all arrows being bidifferential homomorphisms and the vertical arrows being in addition lifting extensions. This says precisely that we have compatible B -variety structures on the base extensions X_L and Y_L such that ϕ_L is a B -morphism.

Now, fix $\alpha = (y_1, \dots, y_m) \in Y_L(L)$ such that y_1, \dots, y_m generate $k[Y]$. So α is a generic point of Y_L over k . Theorem 3.6 also tells us that the graph of ϕ restricted to $X \times Y_0$ is a B -subvariety. So the ideal $I \subseteq k[X] \otimes_k k[Y_0]$ generated by

$$\{\phi^* y_j \otimes 1 - 1 \otimes y_j : j = 1, \dots, m\}$$

is a bidifferential ideal. Hence the extension $I \cdot L[X_L]$ is a bidifferential ideal, and this is precisely the vanishing ideal of the fibre $\phi^{-1}(\alpha)$ in $X_L(L)$. On the other hand, the contraction of $I \cdot L[X_L]$ to $L[Y_L]$, which is precisely the vanishing ideal of $\alpha \in Y_L(L)$, is also bidifferential (being the contraction of a bidifferential ideal under a bidifferential extension). So α is a B -point and the fibre is a B -subvariety. \square

Remark 3.10. Applying the theorem to the case when $X = Y$ and $\phi = \text{id}$ we see that every B -variety has a *generic B -point* over k ; namely a compatible base extension to the function field and a B -point of this extension that is (Zariski) generic over k .

We can iterate this to get a “universal” base extension for $\phi : X \rightarrow Y$.

Corollary 3.11. *Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of absolutely irreducible B -varieties over k . There is a bidifferential field extension $k \subseteq L$ and compatible base extensions X_L and Y_L so that $\phi_L : X_L \rightarrow Y_L$ is a B -morphism, and for every finitely generated subextension field, say $k \subseteq F \subseteq L$, there is a B -point $\alpha \in Y_L^\sharp(L)$ that is generic over F and such that the fibre $\phi_L^{-1}(\alpha)$ is a B -subvariety.*

Proof. We construct a chain of bidifferential field extensions

$$k = L_0 \subseteq L_1 \subseteq L_2 \subseteq \dots$$

with compatible B -variety structures on X_{L_i}, Y_{L_i} such that $\phi_{L_i} : X_{L_i} \rightarrow Y_{L_i}$ are B -morphisms, and $Y_{L_i}(L_i)$ has a B -point that is generic over L_{i-1} and such that the fibre above that point is a B -subvariety of X_{L_i} . Indeed, we go from L_i to $L_{i+1} := L_i(Y_{L_i})$ by Theorem 3.9. Then $L := \bigcup_{i < \omega} L_i$ works. \square

Definition 3.12 (Universal base extension). Given $\phi : X \rightarrow Y$, a dominant B -morphism of absolutely irreducible B -varieties over k , we call the B -morphism $\phi_L : X_L \rightarrow Y_L$ given by Corollary 3.11 a *universal base extension* for $\phi : X \rightarrow Y$.

4. THE BIDIFFERENTIAL DIXMIER-MOEGLIN EQUIVALENCE PROBLEM

We now extend the Poisson Dixmier-Moeglin equivalence problem to this setting of bidifferential algebra, and make some observations about it.

Fix a bidifferential field $(k, \{\cdot, \cdot\})$ of characteristic zero, and $(R, \{\cdot, \cdot\})$ an affine bidifferential k -algebra. Recall that here affine means that R is an integral domain which is finitely generated as a k -algebra.

Given an ideal I of R , it is not hard to check that there is a bidifferential ideal $J \subseteq I$ that contains all other bidifferential ideals of I (this follows from the fact that if J_1 and J_2 are bidifferential ideals of R then $J_1 + J_2$ is bidifferential). We call J the *bidifferential core* of I .

We denote by $\text{Spec}_B(R)$ the space of prime bidifferential ideals of R . We introduce the following three conditions, inspired by the Poisson Dixmier-Moeglin equivalence problem:

Definition 4.1. Let $P \in \text{Spec}_B(R)$.

- (a) P is *B-locally-closed* if $P \neq \bigcap_{P \subsetneq Q \in \text{Spec}_B(R)} Q$.
- (b) P is *B-primitive* if it is the bidifferential core of a maximal ideal.
- (c) P is *B-rational* if the constants of $\text{Frac}(R/P)$ are contained in k^{alg} .

The last of these may require some explanation. As P is a bidifferential ideal we have an induced biderivation on R/P , which by Corollary 2.9 extends uniquely to the fraction field. Recall that by the *constants* of $\text{Frac}(R/P)$ we mean those $\alpha \in \text{Frac}(R/P)$ such that the associated hamiltonians $\{\alpha, \cdot\}$ and $\{\cdot, \alpha\}$ are identically zero on $\text{Frac}(R/P)$.

Note that when $\{\cdot, \cdot\}$ is a Poisson bracket on R , these conditions coincide precisely with Poisson locally closed, Poisson primitive, and Poisson rational, considered by, for example, Brown and Gordon [5]. They are also naturally analogous to the differential version studied in [2].

We begin by showing that the expected implications hold also in this setting. We include proofs for the sake of completeness, though there is no significant difficulty in passing from the Poisson case to the general bidifferential one.

But first a lemma.

Lemma 4.2. *The radical of a bidifferential ideal is bidifferential. If a radical differential ideal is the intersection of finitely many prime ideals none of which contains the other, then these prime ideals are bidifferential.*

Proof. This follows by the corresponding facts about derivations. Namely, that in a differential ring the radical of a differential ideal is differential, and if a radical differential ideal is the intersection of finitely many primes, no one containing the other, then those primes are all also differential. See [9, §0.9 and §1.2]. Now just apply this to each of the hamiltonians $\{r, \cdot\}$ and $\{\cdot, r\}$ for each $r \in R$. \square

Proposition 4.3. *For any prime bidifferential ideal we have*

$$B\text{-locally-closed} \implies B\text{-primitive},$$

and, if k is contained in the constants of $(R, \{\cdot, \cdot\})$ then we also have

$$B\text{-primitive} \implies B\text{-rational}.$$

Proof. Working in the quotient ring, it suffices to consider the case of the prime bidifferential ideal (0) .

Assume (0) is B -locally-closed. Let

$$J = \bigcap_{(0) \neq Q \in \text{Spec}_B(R)} Q.$$

Then J is a nonzero radical bidifferential ideal. Let \mathfrak{m} be a maximal ideal of R not containing J . We claim that (0) is the bidifferential core of \mathfrak{m} (which shows that (0) is B -primitive). For a contradiction, suppose I is a nonzero bidifferential ideal contained in \mathfrak{m} . By Lemma 4.2, the radical of I is bidifferential and its prime components are bidifferential. One of these components must be contained in \mathfrak{m} . However, all these components contain J , contradicting the fact that $J \not\subseteq \mathfrak{m}$.

Now assume that k is contained in the constants of $(R, \{\cdot, \cdot\})$. Suppose that (0) is B -primitive. Namely, there is a maximal ideal \mathfrak{m} with bidifferential core (0) . Let r be a constant in $(\text{Frac}(R), \{\cdot, \cdot\})$. We need to show that $r \in k^{\text{alg}}$.

We claim that there are $a, b \in R$ with $r = a/b$ and $b \notin \mathfrak{m}$. Assume on the contrary that for every representation of $r = a/b$ we have $b \in \mathfrak{m}$. Our goal is to deduce a contradiction to (0) being the differential core of \mathfrak{m} by showing that the bidifferential ideal generated by b is contained in \mathfrak{m} . For this it suffices to show that if you iteratively apply hamiltonians to b then you stay in \mathfrak{m} . By symmetry we can consider only hamiltonians of the form $\{c, \cdot\}$ with $c \in R$. Now, since r is a constant, we get

$$0 = \{c, r\} = \{c, a/b\} = \frac{\{c, a\}b - a\{c, b\}}{b^2}.$$

Then, either $\{c, b\} = 0$ or $r = \frac{a}{b} = \frac{\{c, a\}}{\{c, b\}}$. In either case, $\{c, b\} \in \mathfrak{m}$. Moreover, if $\{c, b\} = 0$ then of course further applications of hamiltonians will stay in \mathfrak{m} , and if $\{c, b\} \neq 0$ then $r = \frac{\{c, a\}}{\{c, b\}}$ and so can repeat the argument with another hamiltonian. hence, applying iterated hamiltonians to b stays in \mathfrak{m} , as desired.

We can thus write $r = a/b$ such that $b \notin \mathfrak{m}$. Work in the localisation R_b on which we can uniquely extend the biderivation by Theorem 2.8. Since $b \notin \mathfrak{m}$, we have that $\mathfrak{m}R_b$ is a maximal ideal of R_b , and so $R_b/\mathfrak{m}R_b$ is an algebraic extension of k . It follows that there is a nonconstant polynomial $P(t) \in k[t]$ such that $P(r) \in \mathfrak{m}R_b$. Note that the constants of $(R_b, \{\cdot, \cdot\})$ contain k and r , and hence $\{s, P(r)\} = \{P(r), s\} = 0$ for all $s \in R_b$. It follows that the bidifferential ideal generated by $P(r)$ in $(R_b, \{\cdot, \cdot\})$ is just the principal ideal $P(r)R_b \subseteq \mathfrak{m}R_b$. But this implies that the contraction $P(r)R_b \cap R$ is a bidifferential ideal of $(R, \{\cdot, \cdot\})$ contained in \mathfrak{m} . As the bidifferential core of \mathfrak{m} is trivial, we must have $P(r) = 0$, proving that $r \in k^{\text{alg}}$. \square

We are therefore lead to pose:

Problem 4.4 (Bidifferential Dixmier-Moeglin Equivalence – BDME). For which affine bidifferential k -algebras is it the case that every B -rational prime bidifferential ideal is B -locally closed?

Any progress on this problem would of course also be progress on the Poisson DME problem. It is known that the PDME does not hold for all affine Poisson k -algebras, see [1]. It is conjectured that the PDME does hold for Poisson-Hopf k -algebras, and some partial results have been obtained, see [10]. The differential

analogue of the DME has also been studied, and various results, including its failure in general and its truth in the Hopf case, have been established, see [2].

It may be worth pointing out that the following weakening of “rational implies locally closed” was shown for affine Poisson algebras in [1, Theorem 7.1], but the proof goes through more or less verbatim in the general bidifferential setting:

Fact 4.5. *Suppose $(R, \{\cdot, \cdot\})$ is an affine bidifferential k -algebra with constants extending k . If $P \in \text{Spec}_B(R)$ is rational then there are only finitely many prime bidifferential ideals containing P of height equal to $\text{ht } P + 1$.*

Remark 4.6. As a consequence of Fact 4.5, and following the ideas of [1, Lemma 7.2], one can show that the BDME holds of affine bidifferential rings of Krull dimension at most 2. Moreover, as long there exists $r \in R \setminus k^{\text{alg}}$ such that $\{r, r\} = 0$ (this condition holds for all r when $\{\cdot, \cdot\}$ is skew-symmetric), then the BDME follows in Krull dimension at most 3.

4.1. An image and generic fibre approach. We conclude by illustrating how the constructions of Section 3 may be of use in this problem, and why passing to general bidifferential algebras over possibly nonconstant base fields may be useful even if one is primarily interested in the Poisson case.

First, to situate the problem in a geometric setting, let us say for short that an irreducible B -variety, X , over k , is B -locally-closed if in $(k[X], \{\cdot, \cdot\})$ the zero ideal is B -locally closed. Geometrically, this is the property that the union of its proper B -subvarieties is not Zariski dense.

Lemma 4.7. *An irreducible B -variety is B -locally-closed if and only if it has only finitely many maximal proper irreducible B -subvarieties.*

Proof. It is clear that if X has only finitely many maximal proper irreducible B -subvarieties then X must be B -locally closed, as the union of these would be a proper Zariski closed set containing all the proper B -subvarieties.

Suppose X is B -locally-closed and let

$$I := \bigcap \{Q \subseteq k[X] : Q \neq 0 \text{ prime bidifferential ideal}\}.$$

Let P_1, \dots, P_ℓ be the minimal associated prime ideals of I , and denote by Z_i the closed subvariety of X defined by P_i . By Lemma 4.2, we get that each Z_i is a B -subvariety. By definition, every proper irreducible B -subvariety of X is contained in some Z_i . Since I , and hence each P_i , is nonzero, we have that Z_1, \dots, Z_ℓ are the maximal irreducible proper B -subvarieties of X . \square

The following gives an “image and generic fibre” characterisation of B -locally closed.

Proposition 4.8. *Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of irreducible B -varieties over k . Assume Y is B -locally-closed. Then the following are equivalent:*

- (i) X is B -locally-closed
- (ii) The generic B -fibre of ϕ over $k(Y)$ is B -locally-closed.

Proof. Let $L := k(Y)$ equipped with the induced bidifferential field structure. Fix compatible B -variety structures on X_L and Y_L such that $\phi_L : X_L \rightarrow Y_L$ is a

B -morphism, and fix $\alpha \in Y_L^\sharp(L)$ generic over k such that $X_\alpha := \phi_L^{-1}(\alpha)$ is a B -subvariety of X_L . This is possible because of Theorem 3.9.

Note that $k(\alpha) = k(Y) = L$. The generic B -fibre referred to in (ii) is X_α .

We will show, first of all, that there is a bijective containment preserving correspondence between

$$\mathcal{X} := \{\text{irreducible } B\text{-subvarieties of } X \text{ over } k \text{ that map dominantly onto } Y\}$$

and

$$\mathcal{X}_\alpha := \{\text{irreducible } B\text{-subvarieties of } X_\alpha \text{ over } L\}$$

given by

$$X' \mapsto X'_\alpha := X'_L \cap X_\alpha.$$

That this is true if we replace “ B -subvariety” by “subvariety” follows from the fact that α is generic in Y over k . So what we are claiming is that X' is a B -subvariety if and only if X'_α is. Indeed, suppose $X' \in \mathcal{X}$. Then, as the base extension X_L has a compatible B -variety structure, X'_L is a B -subvariety of X_L . Hence, so is $X'_L \cap X_\alpha = X'_\alpha$. Conversely, suppose $Z \in \mathcal{X}_\alpha$. Let $X' \subseteq X$ be the subvariety over k with $X'_\alpha = Z$. Note that $I(X') = I(Z) \cap k[X]$. So, as $k[X]$ is a bidifferential subring of $L[X_L]$, we have that X' is a B -subvariety. We thus have a bijective correspondence as claimed.

Assume now that (i) holds. Note that the maximal elements of \mathcal{X} are also maximal among *all* irreducible B -subvarieties of X (over k), as any B -subvariety containing an element of \mathcal{X} will also map dominantly onto Y . Hence, by (i), \mathcal{X} has only finitely many maximal elements. By the bijective correspondence, \mathcal{X}_α has only finitely many maximal elements, thus establishing (ii).

Now, suppose (ii) holds so that \mathcal{X}_α has only finitely many maximal elements. The bijective correspondence gives us that \mathcal{X} has only finitely many maximal elements. Let X_0 be the union of these. So every proper irreducible B -subvariety of X that maps dominantly onto Y is contained in X_0 . Let Y_0 be a proper algebraic subvariety of Y that contains all proper B -subvarieties of Y ; it exists by our assumption on Y . Note that the Zariski closure of the image of a B -subvariety under a B -morphism is a B -subvariety; this is because the intersection of a bidifferential ideal with a bidifferential subring is a bidifferential ideal of the subring. Hence, those B -subvarieties of X that do not project dominantly onto Y must land in Y_0 . It follows that every proper B -subvariety of X lands in $X_0 \cup \phi^{-1}(Y_0)$. The latter being a proper subvariety of X by irreducibility, this proves (i). \square

Similarly, we can say that an irreducible B -variety over k is *B -rational* if the zero ideal in $k[X]$ is B -rational. Geometrically, this means that any rational function $f \in k(X)$ such that $\{f, -\} \equiv 0$ on $k(X)$ must be algebraic over k . It is even easier to see that this also has an “image and generic fibre” characterisation:

Proposition 4.9. *Suppose $\phi : X \rightarrow Y$ is a dominant B -morphism of irreducible B -varieties over k . Assume that Y is B -rational. Then the following are equivalent:*

- (i) X is B -rational.
- (ii) The generic B -fibre of ϕ over $k(Y)$ is B -rational.

Proof. As before, let $L := k(Y)$ and use Theorem 3.9 to fix compatible B -variety structures on X_L and Y_L such that $\phi_L : X_L \rightarrow Y_L$ is a B -morphism, and fix

$\alpha \in Y_L^\sharp(L)$ generic over k such that $X_\alpha := \phi_L^{-1}(\alpha)$ is a B -subvariety of X_L . So $L = k(\alpha)$ and X_α is our generic B -fibre referred to in (ii).

Note that $k(X) = L(X_\alpha)$. This immediately shows that (i) implies (ii). For the converse, assume (ii) and let $f \in k(X)$ be such that $\{f, -\} \equiv 0$. Then we must have $f \in L$. But $L = k(Y)$ and so our assumption on Y implies that $f \in k^{\text{alg}}$, establishing (ii). \square

Propositions 4.8 and 4.9 suggest an inductive approach to Problem 4.4.

It is also worth observing that, even if we start with a Poisson variety X , the B -variety structure we are considering on the generic fibres X_α in these propositions will not be Poisson and will be over nonconstant base fields.

APPENDIX A. EXTENDING DERIVATIONS

We review some basic facts about extending derivations. We could not find good references because we work in the slightly more general, and somewhat unusual, multisorted setting. That is, we work with the collection of pairs (R, M) where R is a (commutative unitary) ring and M is an R -module. Given such (R, M) , we consider M -valued derivations on R , namely additive maps $d : R \rightarrow M$ that satisfy the Leibniz rule $d(rs) = d(r)s + rd(s)$. By a *morphism* $(R, M) \rightarrow (R', M')$, we mean a pair (ϕ, ψ) where $\phi : R \rightarrow R'$ is a ring homomorphism and $\psi : M \rightarrow M'$ is a group homomorphism that is R -linear in the sense that $\psi(rx) = \phi(r)\psi(x)$, for all $r \in R$ and $x \in M$. Given derivations $d : R \rightarrow M$ and $d' : R' \rightarrow M'$, a morphism $(R, M) \rightarrow (R', M')$ is *differential* if $\psi(dr) = d'r$. In that case, we say that $d' : R' \rightarrow M'$ *extends* $d : R \rightarrow M$.

Definition A.1 (Differential triviality). An A -algebra R is said to be *differentially trivial* if R admits no nontrivial A -linear derivations (with values in any R -module).

Differential triviality ensures that any derivation on A has at most one extension to R . So, for example, (separably) algebraic field extensions are easily seen to be differentially trivial. Moreover any derivation on a field lifts to algebraic extensions, so we have existence and uniqueness. If you consider integral extensions of domains instead of algebraic extensions of fields, one does not quite have differential triviality. Nevertheless, the following approximation holds:

Fact A.2. *Suppose $A \subseteq B$ is an extension of integral domains and $b \in B$ is algebraic over $\text{Frac}(A)$ with the minimal polynomial P of b over $\text{Frac}(A)$ having coefficients in A . Suppose S is a B -algebra. Then any derivation $d : A \rightarrow S$ extends uniquely to a derivation $A[b] \rightarrow S_f$ where $f := \frac{dP}{dt}(b)$.*

Proof. The statement is meant to be understood with respect to the natural morphism $(A, S) \rightarrow (A[b], S_f)$.

Uniqueness follows from the fact that $P(b) = 0$ forces $d(b) = -\frac{P^d(b)}{f}$, where $P^d \in S[t]$ is obtained by applying d to the coefficients of P .

For existence, first we make S_f an algebra over the polynomial ring $A[t]$ by the evaluation map $t \mapsto b$. Next we extend $d : A \rightarrow S$ to a derivation $d : A[t] \rightarrow S_f$ by setting $dt := -\frac{P^d(b)}{f}$. Note that the transcendence of t means this is always possible, and that for any $Q \in A[t]$ we have $dQ = \frac{dQ}{dt}(b)dt + Q^d(b)$. In particular, $dP = 0$. Let $I \subseteq A[t]$ be the ideal of polynomials vanishing at b . As $P \in A[t]$ is

monic, and of minimal degree satisfying $P(b) = 0$, the division algorithm for monic divisors in $A[t]$ implies that $I = (P)$. It follows that $dQ = 0$ for all $Q \in I$. So we get the desired induced S_f -valued derivation on $A[b] = A[t]/I$. \square

Remark A.3. The above fact only has content when f is not nilpotent, as otherwise the localisation S_f is the trivial ring. In fact, it is usually applied when b is separably algebraic over $\text{Frac}(A)$ and S has no B -torsion, in which case f is not nilpotent.

Derivations amalgamate nicely with respect to tensor products:

Fact A.4. *Suppose we have a commuting diagram of ring homomorphisms*

$$\begin{array}{ccc} & R_1 & \\ & \nearrow & \searrow \\ A & & S \\ & \searrow & \nearrow \\ & R_2 & \end{array}$$

and derivations $d_1 : R_1 \rightarrow S$ and $d_2 : R_2 \rightarrow S$ that agree on A . Then there is a unique derivation $d : R_1 \otimes_A R_2 \rightarrow S$ extending both d_1 and d_2 .

Proof. Define $D : R_1 \times R_2 \rightarrow S$ by $D(x, y) = (d_1 x)y + x(d_2 y)$. This is \mathbb{Z} -bilinear and hence induces an additive map $d : R_1 \otimes_{\mathbb{Z}} R_2 \rightarrow S$, that is easily seen to be a derivation. Moreover, for each $a \in A$ we have

$$d(a \otimes 1 - 1 \otimes a) = (d_1 a)1 + a(d_2 1) - (d_1 1)a - 1(d_2 a) = d_1 a - d_2 a = 0$$

since d_1 and d_2 agree on A . So we obtain an induced derivation $d : R_1 \otimes_A R_2 \rightarrow S$. The Leibniz rule ensures uniqueness. \square

REFERENCES

- [1] J. Bell, S. Launois, O. León Sánchez, and R. Moosa. Poisson algebras via model theory and differential-algebraic geometry. *J. Eur. Math. Soc. (JEMS)*, 19(7):2019–2049, 2017.
- [2] J. Bell, O. León Sánchez, and R. Moosa. D -groups and the Dixmier-Moeglin equivalence. *Algebra Number Theory*, 12(2):343–378, 2018.
- [3] M. Bresar. Jordan g, h -derivations on tensor products of algebras. *Linear and Multilinear Algebra*, 64(11):2199–2207, 2016.
- [4] M. Bresar, W. S. Martindale, and C. R. Miers. Centralizing maps in prime rings with involution. *J. Algebra*, 161:342–357, 1993.
- [5] K. A. Brown and I. Gordon. Poisson orders, symplectic reflection algebras and representation theory. *J. Reine Angew. Math.*, 559:193–216, 2003.
- [6] A. Buium. *Differential algebraic groups of finite dimension*, volume 1506 of *Lecture Note in Mathematics*. Springer-Verlag, 1992.
- [7] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [8] D. Eremita. Biderivations on tensor product of algebras. *Communications in Algebra*, 46(4):1722–1726, 2018.
- [9] E.R. Kolchin. *Differential algebra and algebraic groups*, volume 54 of *Pure and Applied Mathematics*. Academic Press, New York-London, 1973.
- [10] S. Launois and O. León Sánchez. On the Dixmier-Moeglin equivalence for Poisson-Hopf algebras. *Adv. Math.*, 346:48–69, 2019.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, OXFORD ROAD, MANCHESTER,
UNITED KINGDOM M13 9PL

Email address: `omar.sanchez@manchester.ac.uk`

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVENUE
WEST, WATERLOO, ONTARIO, CANADA N2L 3G1

Email address: `rmoosa@uwaterloo.ca`